

DRUG FACTS

Active ingredient(s)

Purpose

Use(s)

temporarily relieves common cold/flu symptoms:

- nasal congestion (Daytime only)
- sore throat
- cough due to minor throat and bronchial irritation
- headache
- minor aches and pains
- fever
- runny nose and sneezing (Nighttime only)

Warnings

Liver warning: These products contain acetaminophen. Severe liver damage may occur if:

- adult takes more than 4 doses in 24 hours (Nighttime) or more than 6 doses in 24 hours (Daytime), which is the maximum daily amount
- child takes more than 5 doses in 24 hours (Daytime)
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using these products

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

Ask a doctor before use if you have

- high blood pressure (Daytime only)
- heart disease (Daytime only)
- diabetes (Daytime only)
- thyroid disease (Daytime only)
- trouble urinating due to an enlarged prostate gland
- liver disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma or emphysema
- a breathing problem such as emphysema or chronic bronchitis (Nighttime only)
- glaucoma (Nighttime only)
- a sodium-restricted diet (Nighttime only)

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- do not use more than directed
- In addition, when using Nighttime:
- avoid alcoholic drinks
 - marked drowsiness may occur
 - excitability may occur, especially in children
 - alcohol, sedatives and tranquilizers may increase drowsiness
 - be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless (Daytime only)
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as recommended – see Liver warning
- take Nighttime or Daytime
- use dose cup

Nighttime Liquid

- do not exceed 4 doses per 24 hours

adults and children 12 years and over	30 mL (2 TBSP) every 6 hours
children 4 to under 12 years	ask a doctor
children under 4 years	do not use

Daytime Liquid

- do not exceed 5 doses (children) or 6 doses (adults) per 24 hours

adults and children 12 years and over	30 mL (2 TBSP) every 4 hours
children 6 to under 12 years	15 mL (1 TBSP) every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

- when using other Daytime or Nighttime products, carefully read each label to insure correct dosing

Other information

- each TBSP of Nighttime contains: sodium 18 mg
- each TBSP of Daytime contains: sodium 7 mg

- store at 20°-25°C (68°-77°F)

Inactive ingredients

Nighttime alcohol, anhydrous citric acid, D&C yellow no. 10, FD&C green no. 3, FD&C yellow no. 6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Daytime butylated hydroxyanisole, edetate disodium, FD&C yellow no. 6, flavors, glycerin, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum